

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/312911440>

Natural Compounds Targeting Transforming Growth Factor- β

Article in *Electronic Journal of Biology* · December 2016

CITATIONS

0

READS

21

3 authors:



[Pushendra Singh](#)

National Institute of Pathology

23 PUBLICATIONS 156 CITATIONS

[SEE PROFILE](#)



[Felix Bast](#)

Central University of Punjab

65 PUBLICATIONS 214 CITATIONS

[SEE PROFILE](#)



[Ravi Shankar Singh](#)

Sanjay Gandhi Post Graduate Institute of Me...

10 PUBLICATIONS 46 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Latitudinal gradient in species distribution of phytoplankton in Southern Ocean through metagenomics [View project](#)



DNA Barcode-based assessment of the biodiversity of ice algae, snow algae and lichen-photobionts of East Antarctica [View project](#)

All content following this page was uploaded by [Ravi Shankar Singh](#) on 02 March 2017.

The user has requested enhancement of the downloaded file.

Natural Compounds Targeting Transforming Growth Factor- β : *In Silico* and *In Vitro* Study

Pushpendra Singh^{1,*}, Felix Bast¹, Ravi Shankar Singh²

¹ Centre for Biosciences, School of Basic and Applied Sciences, Central University of Punjab,
Bathinda, Punjab, India;

² Department of Molecular Medicine & Biotechnology, Sanjay Gandhi Post Graduate Institute of
Medical Sciences, Lucknow, India.

*Corresponding author. Tel: 9569330844; Email: pushsingh02@gmail.com

Citation: Singh P, Bast F, Singh RS. Natural Compounds Targeting Transforming Growth Factor- β : *In Silico* and *In Vitro* Study. Electronic J Biol, 13:1

Received: September 21, 2016; **Accepted:** November 29, 2016; **Published:** December 06, 2016

Research Article

Abstract

Inhibition of the tumor-promoting effects of transforming growth factor beta receptor (TGF β R) in carcinogenesis provides a better therapeutic intervention. Various natural compounds, inhibitors of TGF β R have been used for *in vitro* and *in vivo* anticancer study. Although very few TGF β R inhibitors are now intensifying in preclinical studies. In this study our aim to investigate TGF β R1, TGF β R2 and TAK1 inhibitor by using molecular docking and *in vitro* study. Our result revealed that some compounds have better docking energy. Moreover, the effect of two lead molecules epigallocatechin gallate (EGCG) and myricetin on the mRNA expression of TGF β R1 was reported after the 48 hrs treatments in HepG2 and PC3 cancer cell lines. The RT-PCR showed that compound EGCG and myricetin reduced the mRNA expression of TGF β R1 at 80 μ M concentration. This molecular docking study provides a better understanding of binding of compounds to the active site of proteins and to summarize the various binding energy, hydrophobic, hydrogen, an electrostatic bond that are decisive for the protein-ligand interactions. Further experimental work will be required for validation of our results.

Keywords: Natural compounds; Cancer; Maestro 9.6; Molecular docking; *In vitro*

1. Introduction

The Transforming Growth Factor β (TGF β) family have a group of multifunctional regulatory proteins that balance a vast variety of physiological functions including cell proliferation, cell cycle arrest and apoptosis. TGF β R1 53kDa protein is also known as activin encoded by TGF β R1 human gene. TGF β activated protein kinase 1 (TAK1) was acknowledged for protein kinase action that invigorating by TGF β signaling transduction [1]. Furthermore, it is also demonstrated that TAK1 cytokine play a significant role in inflammation. TGF β initiates signals by

binding to the TGF β R2 and stabilizes the heteromeric complex with the TGF β R1. As a result, TGF β R1 is trans-phosphorylated and activated by TGF β R2 [2,3]. The activated TGF β R1 then propagates the signals through interaction with receptor-associated Smads [4]. Deregulations of TGF β signaling pathway responsible for cancer initiation and progression and interrupting the tumor promoter properties of TGF β signaling would be an attractive therapeutic strategy [5-7]. Monoclonal antibodies and small molecule inhibitors that target TGF β signaling are the most notable strategy have been used in preclinical studies, but they did not prove much promise as anticancer drugs owing to multifaceted roles of TGF β signaling [8,9].

In recent years, natural compounds have been investigated intensely in the clinical trial of ovarian, breast, cervical, pancreatic and prostate cancers [10]. A number of propitious agents are in clinical development based on discriminatory activity against anti-cancer molecular targets currently being developed. In this context, we studied TGF β signaling inhibitors (TGF β R1, TGF β R2, and TAK1) from natural sources using *in silico* and *in vitro* approaches.

2. Material and Methods

2.1 Selection of ligands and protein molecules

In silico, molecular docking protocol and standard setting of parameter adapted from our published literature [11-16]. Marine and natural compounds selected as ligand molecules that had been reported as an anticancer activity in the published literature [17-29]. The X-ray crystal structure of TGF β R1 (PDB; 2X7O), TGF β R2 (PDB; 1KS6), and TAK1 (PDB; 4L3P) kinase protein retrieved from the Protein Data Bank (PDB) (Figure 1).

2.2 Preparation of ligands molecules

Input ligand molecules were prepared using ligprep wizard applications where the addition of hydrogen atoms, 2D to 3D conversion followed by optimized




Figure 1. Workflow of study design.

potential for liquid simulations (OPLS_2005) force field was applied [30-32]. Finally, ten conformations for each ligand were generated and performed molecular docking.

2.3 Preparation of protein molecules

Protein molecules were prepared using respective wizard applications (standard methods) where changes such as the addition of hydrogen atoms, fixing of the charges and orientation of groups, assigning bond orders, creation of disulphide bonds, was done into the PDB structure. After the completion of ligands and protein preparation, a receptor-grid file was generated.

2.4 GLIDE (Grid-based Ligand Docking with Energetics) molecular docking

The active site (binding pocket) and functional residues of protein were identified and characterized by GLIDE from Schrodinger package. GLIDE-XP Molecular docking studies using the selected ligand molecules were conducted using Maestro 9.6 [33-35]. Each of these selected compounds was docked into target protein molecules and a compound possessing the lowest energy was selected. After the configuration of receptor-grid file, flexible ligands with rigid receptor molecular docking were performed. The concluding energy evaluation was done on the basis of G-score.

2.5 Absorption, distribution, metabolism, excretion and toxicity (ADME/T) studies

ADME properties such as poor compound solubility, gastric emptying time and lack of ability to permeate

the intestinal wall can all diminish the extent to which a drug is absorbed after oral administration. Therefore, *in silico* ADME/T predictive tools are accommodating approach that could abolish unfortunate compounds, before invested valuable time and money in primary testing of compounds. QikProp application of Maestro 9.6 predicts properties such as logBB, overall CNS activity, Caco-2, and MDCK cell permeability and logK_{hsa} for human serum albumin binding, etc. [30-36].

2.6 Reagents and cell lines culture

DMEM media, 1% penicillin, streptomycin and Fetal Bovine Serum (FBS) procured from Invitrogen, and Ham's F-12 media from HiMedia. EGCG and myricetin purchased from MP Biomedicals Pvt. Ltd. and dissolved in Dimethyl Sulfoxide (DMSO) as a 20 mM stock solution. We used final DMSO concentration in culture medium were 0.25% (vol/vol). Hepatocellular carcinoma (HepG2) and Prostate Cancer (PC3), cell lines were procured from NCCS, Pune, India. Cells were grown in phenol red media containing 10% FBS and 1% penicillin and streptomycin and have to grow in a 37°C incubator with 5% CO₂.

2.7 Total RNA isolation, cDNA synthesis and quantitative RT-PCR

One million cells/well were plated into the six-well culture plate media supplemented with 10% FBS and 1% penicillin/streptomycin then incubated at 37°C overnight with 5% CO₂. Cells were exposed to 80 μM EGCG and Myricetin for 48 h. After 48 h total RNA was extracted using Tri reagent (Life Technologies, Gaithersburg, MD) according to manufacturer's

Table 2: Lowest binding energy for the ligand- TGF β R1 (PDB; 2X7O) and TGF β R2 (PDB; 4XJJ) protein interactions.

Ligand type	Compounds ID	G-Score	Lipophilic Ewdw	HBond	Electro	Protein-ligands interactions
	SD 208	-9.95	-5.15	-0.7	-0.35	Hid283
	R 268712	-8.04	-5.5	-1.2	-0.9	Lys232, Ser287, Asp290
	CID4521392	-7.87	-5.16	-0.81	-0.25	Hid283, Lys335
TGF β R	A 83-01	-7.33	-6.89	-0.05	-0.58	Lys337
inhibitors	GW 788388	-7.29	-6.3	-0.35	-0.42	Lys232
(Controls)	449054	-7.23	-4.73	-0.32	-0.36	Lys232, Hid283
	LY 364947	-7.07	-5.09	-0.17	-0.32	Lys232
	D 4476	-7.05	-5.66	-0.88	-0.29	Lys232, Glu245, Hid183
	CID11655119	-7.03	-7.29	0	-0.34	
	SB 525334	-6.46	-4.8	-0.47	-0.54	Lys232, Hid183
	CID65064	-13.05	-6.2	-5.13	-1.36	Leu278, Hid183, Lys337
	CID72277	-11.6	-4.67	-4.12	-1.22	Lys232, Glu245, Hid183, Asp351
	CID5281672	-11.36	-5.02	-3.96	-0.99	Lys232, Glu245, Hid283
TGF β R1	CID9892144	-11.14	-6.83	-2.8	-0.54	Lys213, Lys232, Hid183, Ser287
	CID72276	-11.09	-4.72	-3.46	-1.12	Lys232, Glu245, Hid183, Asp351
	CID72281	-10.94	-4.97	-2.94	-0.98	Lys232, Glu245, Hid183, Asp351
	CID5280343	-10.93	-4.98	-3.53	-0.99	Lys232, Glu245, Hid183
	CID9064	-10.58	-4.69	-2.96	-1.13	Lys232, Glu245, Hid183, Asp351
	CID6445533	-10.47	-5.29	-3.06	-1.18	Arg215, Leu278, Asp281, Asp351
	CID5280445	-10.42	-4.94	-3.04	-1.01	Lys232, Glu245, Hid183
	CID44259	-10.3	-5.71	-1.33	-0.52	Asp281, Hid283
	CID5281614	-9.98	-4.88	-2.51	-0.82	Asp281, Glu245, Asp351
	CID72276	-8.03	-3.69	-3.51	-0.77	Gln41, Trp65, Met100, Phe111, Phe126
	CID5280343	-7.81	-3.83	-3.03	-0.82	Gln41, Trp65, Met100, Phe111, Phe126
	CID9064	-7.71	-3.71	-3.03	-0.8	Gln41, Trp65, Met100, Phe111, Phe126
TGF β R2	CID5281614	-7.57	-3.81	-3.03	-0.76	Trp65, Met100, Phe126
	CID65064	-7.3	-3.29	-3.3	-1.15	Gln41, Lys42, Trp65, Met100, Phe126
	CID42607750	-6.98	-2.87	-3.07	-1.3	Met100, Lys101, Phe126
	CID6400741	-6.42	-3.32	-2.23	-1.42	Gln41, Trp65, Met100, Phe111, Phe126
	CID5281672	-6.36	-2.57	-2.94	-0.92	Gln41, Trp65, Met100, Glu128, Phe126
	CID636888	-6.27	-4.12	-2.29	-0.6	Gln41, Glu119, Phe126
	CID72277	-6.25	-1.99	-3.87	-0.9	Gln41, Trp65, Phe126
	CID5281605	-5.87	-2.2	-2.4	-0.74	Gln41, Trp65, Phe126
	CID439260	-5.78	-1.93	-2.4	-0.85	Gln41, Phe126

involved in hydrophobic interaction at the active site of the protein. Moreover, Additionally, it have shown that amino acids Val42, Ala 46, Glu105 and Ala107, involved in side-chain hydrogen bonding whereas amino acid Lys63 and Asp175 involved in back-bone hydrogen bonding at the active site of TAK1 with myricetin. Additionally amino acid Arg44 and Ala107 of TAK1 form backbone hydrogen bonding with EGCG (Figure 3 and Table 3).

Previously, it was reported that EGCG can inhibit TGF β induced epithelial-mesenchymal transitions via down-regulation of phosphorylated Smad2 and Erk1/2 in non-small cell lung cancer cells [37]. Moreover, EGCG significantly subdues the HSP27 induction prompted by TGF β in a dose-dependent manner. EGCG significantly suppressed the TGF β induced phosphorylation of stress-activated kinases and c-Jun N-terminal kinases [38]. Furthermore, it was noticed that EGCG interacts with TGF β R2 and

inhibits the expression of α -smooth muscle actin via the TGF β -Smad2/3 pathway in human lung fibroblast cells [39,40]. *In vivo* study showed that myricitrin was noticeably ameliorated the expression of TGF β 1 and alpha-smooth muscle actin that led to antioxidant, anti-inflammatory and antifibrotic activity [41]. Furthermore, *in vivo* study in streptozotocin with cadmium-induced nephrotoxic diabetic rats unveil that myricetin up regulates the TGF β 1 and downregulate the peroxisome proliferator-activated receptor alpha proteins expression that led to restraints dyslipidemia and renal mesangial cell proliferation [42].

4.2 ADME/T studies

ADME/T (absorption, distribution, metabolism, elimination and toxicity,) properties of lead molecules were assessed through the Qikprop application of Maestro 9.6. EGCG and myricetin have the best Gscore against TGF β R1, TGF β R2, and TAK1.

Furthermore, EGCG and myricetin have their excellent QPlogPo/w, QPlogHerg K⁺ channels, QPlogBB, QPlogKP, QPlogK HSA values that accomplish the Lipinski's rule of five. However, these compounds do not have excellent QPP Caco, QPP MDCK values and percentage of human oral absorption. Therefore, EGCG and myricetin structural alteration and optimization are required to predict structures that have better QPP Caco, QPP MDCK and rate of human oral absorption activity (Table 4).

4.3 Effect of EGCG and myricetin on TGFβR1 mRNA expression in HepG2 and PC3 cancer cell lines

We investigated whether EGCG and myricetin could decrease the TGFβR1 expression in HepG2 and PC3 cells. After 48 h of treatment with EGCG and myricetin, determine the mRNA expression of TGFβR1 by using RT-PCR. As shown in Figure 4, RT-PCR demonstrates that EGCG have a prospective for forceful inhibition of TGFβR1. The RT-PCR densitometric bands examination showed that EGCG reduced the mRNA expression of TGFβR1

by 15% in HepG2 (*P<0.05) and 70% (**p<0.001) in PC3 cells at 80 μM concentration. Whereas, RT-PCR densitometric bands investigation showed that myricetin reduced the mRNA expression of TGFβR1 by 20% in HepG2 (*P<0.05) and 80% (**p<0.001) in PC3 cells at 80 μM concentrations. TGFβ signaling plays a leading role in the initiation, promotion, and progression of various types of human cancer including prostate, breast lung, and liver cancer. Several lines of evidence suggest that carcinoma cells frequently lose anti-proliferative response to TGFβ. High levels of TGFβ can promote tumor growth in an autocrine and paracrine manner through the stimulation of angiogenesis and metastasis. Blocking the tumor-promoting effects of TGFβ by small molecule inhibitors provides an excellent therapeutic opportunity to improve the treatment of cancer.

5. Conclusion

TGFβ increases the tumor motility, invasion, and metastasis. Inhibition of TGFβ signaling offers a novel approach for the treatment of cancer. Monoclonal




Figure 3. (a) Ribbon presentation of TAK1 (PDB; 4L3P) protein molecule with CID5281672; (b) Protein-ligand interactions profile of 4L3P with CID5281672.

Table 3: Lowest binding energy for the ligand- TAK1 (PDB; 4L3P) protein interactions.

Ligand type	Compounds CID	G-Score	Lipophilic Evdw	H-Bond	Electro	Protein-ligands interactions
TAK1 Inhibitors	CID9863776	-6.82	-3.29	-2.88	-0.44	Ser111, Asn114
	CID57369807	-6.04	-1.97	-3.02	-0.78	Ser111, Asn114, Asn161
Natural Compounds	CID42607750	-12.01	-5.02	-4.48	-1.24	Arg44, Glu105, Ala107, Ser111, Asn114
	CID5281672	-11.67	-3.84	-4.75	-1.69	Val42, Ala 46, Lys63, Glu105, Asp175
	CID6445533	-11.07	-6.55	-2.36	-0.79	Arg44, Glu105, Ala107, Ser111, Asn114
	CID72277	-10.66	-4.62	-3.83	-0.82	Arg44, Glu105, Ala107, Ser111, Asn114
	CID5280445	-10.2	-4.25	-2.95	-1.57	Val42, Ala 46, Lys63, Glu105, Asp175
	CID5280343	-10.12	-3.81	-3.5	-1.53	Val42, Gly43, Lys63, Glu105, Asp175
	CID9892144	-9.96	-6.97	-2.45	-0.58	Ala107, Ser111
	CID5281614	-9.42	-4.35	-3.3	-1.34	Val42, Ala 46, Glu105, Ala107, Asp175
	CID442793	-9.3	-5.04	-2.42	-1.03	Lys63, Glu105, Ala107, Asp175
	CID5281670	-9.08	-5.16	-2.71	-0.78	Lys63, Ala107, Asp175
	CID9064	-8.87	-5.11	-2.46	-0.89	Val42, Arg44, Ala107, Asp175
CID65064	-8.87	-4.14	-4.41	-0.96	Arg44, Asp175	

Table 4: Evaluation of drug-like properties of the lead molecules by Qikprop Maestro.

Compounds	QPlog P _{ow} (-2.0 to 6.5)	QPlog HERG (acceptable range: above -5.0)	QPP Caco (nm/s) <25-poor >500-great	QPlog BB (-3-1.2)	QPP MDCK (nm/s) <25-poor >500- great	QPlog Kp (-8.0 to -0.1)	QPlog Kh _{sa} (Acceptable range: -1.5 to 1.5).	Percentage of human oral absorption; (<25% is poor and >80% is high)
CID65064 (EGCG)	-0.233	-5.558	1.14	-4.177	0.326	-7.415	-0.437	0.684
CID72277	-0.167	-4.684	22.042	-2.338	8.009	-5.482	-0.562	37.049
CID5281672 (myricetin)	-0.276	-4.894	7.673	-2.827	2.56	-6.301	-0.493	28.209
CID9892144	3.838	-5.92	123.998	-1.804	51.813	-3.591	0.723	86.885
CID72276	0.479	-4.852	60.119	-1.858	23.692	-4.599	-0.43	61.589
CID72281	1.782	-5.025	133.02	-1.525	55.9	-4.057	0.007	75.395
CID5280343	0.384	-5.011	20.522	-2.331	7.414	-5.442	-0.348	52.678
CID9064	0.439	-4.845	52.516	-1.913	20.471	-4.716	-0.429	60.305
CID6445533	1.744	-6.907	35.048	-1.764	14.628	-4.913	-0.194	64.802
CID5280445	0.973	-5.14	41.633	-1.975	15.927	-4.849	-0.187	61.625
CID44259	3.765	-6.13	304.782	-0.05	151.525	-4.131	0.787	93.45
CID5281614	0.505	-5.178	45.418	-1.943	17.497	-4.758	-0.364	59.565
CID42607750	-0.607	-5.487	12.878	-3.215	4.48	-5.266	-0.883	17.34
CID5281614	0.458	-4.924	53.31	-1.808	20.806	-4.65	-0.388	60.532
CID442793	3.271	-4.52	506.992	-1.443	237.406	-2.569	0.154	94.512
CID9571127	2.291	-4.098	0.079	-4.908	0.03	-6.406	-0.439	0
CID23663412	3.425	-4.65	8.604	-4.303	3.837	-3.649	-0.49	37.809

QPlog P_{ow} (-2.0 to 6.5) Predicted octanol/water partition coefficient; QPlog Herg (acceptable range: above -5.0) Predicted IC50 value for blockage of HERG K+ channels; QPPCaco (nm/sec) <25-poor >500- great Predicted apparent Caco-2 cell permeability in nm/s. Caco-2 cells is a model for the gut blood barrier; QPlogBB (-3-1.2) Predicted brain/blood partition coefficient; QPPMDCK (nm/s) <25-poor >500- great Predicted apparent MDCK cell permeability in nm/s. MDCK cells are considered to be a good mimic for the blood-brain barrier. QPlogKp- Predicted skin permeability; Q P log Kh_{sa}; Prediction of binding to human serum albumin; (acceptable range: -1.5 to 1.5); Percentage of human oral absorption; (<25% is poor and >80% is high)




Figure 4. Represented effect of EGCG on TGFβR1 mRNA turnover in HepG2 and PC3.

Note: Cells were treated with 80 μM EGCG and myricetin for 48 h of treatment. TGFβR1 mRNA were determined by SYBR® Green based RTPCR using GAPDH as internal control. EGCG and myricetin down-regulated the expression of TGFβR1 mRNA in HepG2 and PC3. Data are s.e.m. (*p<0.05 control vs. EGCG and Myricetin, **p<0.001 control vs. EGCG and myricetin)

antibodies and small compounds used as a TGF β inhibitor but due to cross activity these inhibitors fail in the preclinical trial. Our study showed some compounds has better docking energy such as EGCG and myricetin. Moreover, the RT-PCR showed that compound EGCG and myricetin reduced the mRNA expression of TGF β R1 at 80 μ M concentration. This molecular docking study enhanced understanding of binding of compounds to the active site of proteins and to recapitulate the various binding energy, hydrophobic, hydrogen, an electrostatic bond that are important for the protein-ligand interactions. Further experimental work will be required for validation of our results.

6. Acknowledgement

We would like to thank Vice Chancellor, Central University of Punjab, Bathinda, Punjab, India for supporting this study with infrastructural requirements.

References

- [1] Jakowlew SB. (2006). Transforming growth factor- β in cancer and metastasis. *Cancer and Metastasis Reviews*. **25**: 435-457.
- [2] Ikushima H, Miyazono K. (2010). TGF β signalling: A complex web in cancer progression. *Nature Reviews Cancer*. **10**: 415-424.
- [3] van der Pluijm G. (2011). Epithelial plasticity, cancer stem cells and bone metastasis formation. *Bone*. **48**: 37-43.
- [4] Feng XH, Derynck R. (2005). Specificity and versatility in TGF- β signaling through Smads. *Annu Rev Cell Dev Biol*. **21**: 659-693.
- [5] Kim SS, Shetty K, Katuri V, et al. (2006). TGF- β signaling pathway inactivation and cell cycle deregulation in the development of gastric cancer: Role of the β -spectrin, ELF. *Biochemical and Biophysical Research Communications*. **344**: 1216-1223.
- [6] Xu Y, Pasche B. (2007). TGF- β signaling alterations and susceptibility to colorectal cancer. *Human Molecular Genetics*. **16**: R14-R20.
- [7] Yingling JM, Blanchard KL, Sawyer JS. (2004). Development of TGF- β signalling inhibitors for cancer therapy. *Nature Reviews Drug Discovery*. **3**: 1011-1022.
- [8] Biswas S, Guix M, Rinehart C, et al. (2007). Inhibition of TGF- β with neutralizing antibodies prevents radiation-induced acceleration of metastatic cancer progression. *Journal of Clinical Investigation*. **117**: 1305.
- [9] Hjelmeland MD, Hjelmeland AB, Sathornsumetee S, et al. (2004). SB-431542, a small molecule transforming growth factor- β -receptor antagonist, inhibits human glioma cell line proliferation and motility. *Molecular Cancer Therapeutics*. **3**: 737-745.
- [10] Goel A, Kunnumakkara AB, Aggarwal BB. (2008). Curcumin as "Curecumin": from kitchen to clinic. *Biochemical Pharmacology*. **75**: 787-809.
- [11] Singh P, Bast F. (2015). High-throughput virtual screening, identification and *in vitro* biological evaluation of novel inhibitors of signal transducer and activator of transcription 3. *Medicinal Chemistry Research*. 1-15.
- [12] Singh P, Bast F. (2014a). *In silico* molecular docking study of natural compounds on wild and mutated epidermal growth factor receptor. *Medicinal Chemistry Research*. 1-12.
- [13] Singh P, Bast F. (2014b). Multi-targeted molecular docking study of plant-derived natural products on phosphoinositide-3 kinase pathway components. *Medicinal Chemistry Research*. **23**: 1690-1700.
- [14] Singh P, Bast F. (2015a). Screening and biological evaluation of myricetin as a multiple target inhibitor insulin, epidermal growth factor, and androgen receptor; *in silico* and *in vitro*. *Investigational New Drugs*. **33**: 575-593.
- [15] Singh P, Bast F. (2015b). Screening of multi-targeted natural compounds for receptor tyrosine kinases inhibitors and biological evaluation on cancer cell lines, *in silico* and *in vitro*. *Medical Oncology*. **32**: 1-18.
- [16] Singh P, Singh RS, Rani A, Bast F. (2016). Homology modeling of chemokine CCR7, molecular docking and *in vitro* studies evidenced plausible immunotherapeutic anticancer natural compounds. *Medicinal Chemistry Research*. 1-15.
- [17] Bhanot A, Sharma R, Noolvi MN. (2011). Natural sources as potential anti-cancer agents: A review. *International Journal of Phytomedicine*. **3**: 9-26.
- [18] Cho JY, Park J. (2008). Contribution of natural inhibitors to the understanding of the PI3K/PDK1/PKB pathway in the insulin-mediated intracellular signaling cascade. *International Journal of Molecular Sciences*. **9**: 2217-2230.
- [19] Cragg GM, Newman DJ. (2005). Plants as a source of anti-cancer agents. *Journal of Ethnopharmacology*. **100**: 72-79.
- [20] da Rocha AB, Lopes RM, Schwartzmann G. (2001). Natural products in anticancer therapy. *Current Opinion in Pharmacology*. **1**: 364-369.
- [21] Hillman GG. (2012). Dietary agents in cancer chemoprevention and treatment. *Journal of Oncology*. 1-2.
- [22] Huang W, He T, Chai C, et al. (2012). Triptolide inhibits the proliferation of prostate cancer cells and down-regulates SUMO-specific protease 1 expression. *Public Library of Science One*. **7**: e37693.
- [23] Mayer AM, Gustafson KR. (2004). Marine pharmacology in 2001–2: Anti-tumour and cytotoxic compounds. *European Journal of Cancer*. **40**: 2676-2704.
- [24] Mayer AM, Gustafson KR. (2006). Marine pharmacology in 2003–2004: Anti-tumour and cytotoxic compounds. *European Journal of Cancer*. **42**: 2241-2270.
- [25] Mayer AM, Gustafson KR. (2008). Marine pharmacology in 2005–2006: Anti-tumour and cytotoxic compounds. *European Journal of Cancer*. **44**: 2357-2387.
- [26] Phosrithong N, Ungwitayatorn J. (2010). Molecular docking study on anticancer activity of plant-derived natural products. *Medicinal Chemistry Research*. **19**: 817-835.
- [27] Roell D, Baniahmad A. (2011). The natural compounds atraric acid and N-butylbenzene-sulfonamide as

- antagonists of the human androgen receptor and inhibitors of prostate cancer cell growth. *Molecular and Cellular Endocrinology*. **332**: 1-8.
- [28] Sarkar FH, Li Y. (2006). Using chemopreventive agents to enhance the efficacy of cancer therapy. *Cancer Research*. **66**: 3347-3350.
- [29] Sunil H. (2012). Inhibition studies of naturally occurring terpene based compounds with cyclin-dependent kinase 2 enzyme. *Journal of Computer Science & Systems Biology*. 1-2.
- [30] Jorgensen WL, Duffy EM. (2002). Prediction of drug solubility from structure. *Advanced Drug Delivery Reviews*. **54**: 355-366.
- [31] Jorgensen WL, Maxwell DS, Tirado-Rives J. (1996). Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids. *Journal of American Chemical Society*. **118**: 11225-11236.
- [32] Jorgensen WL, Tirado-Rives J. (1988). The OPLS [optimized potentials for liquid simulations] potential functions for proteins, energy minimizations for crystals of cyclic peptides and crambin. *Journal of American Chemical Society*. **110**: 1657-1666.
- [33] Friesner RA, Banks JL, Murphy RB, et al. (2004). Glide: A new approach for rapid, accurate docking and scoring. 1. Method and assessment of docking accuracy. *Journal of Medicinal Chemistry*. **47**: 1739-1749.
- [34] Friesner RA, Murphy RB, Repasky MP, et al. (2006). Extra precision glide: Docking and scoring incorporating a model of hydrophobic enclosure for protein-ligand complexes. *Journal of Medicinal Chemistry*. **49**: 6177-6196.
- [35] Halgren TA, Murphy RB, Friesner RA, et al. (2004). Glide: A new approach for rapid, accurate docking and scoring. 2. Enrichment factors in database screening. *Journal of Medicinal Chemistry*. **47**: 1750-1759.
- [36] Lu JJ, Crimin K, Goodwin JT, et al. (2004). Influence of molecular flexibility and polar surface area metrics on oral bioavailability in the rat. *Journal of Medicinal Chemistry*. **47**: 6104-6107.
- [37] Liu LC, Tsao TCY, Hsu SR, et al. (2012). EGCG inhibits transforming growth factor- β -mediated epithelial-to-mesenchymal transition via the inhibition of Smad2 and Erk1/2 signaling pathways in non-small cell lung cancer cells. *Journal of Agricultural and Food Chemistry*. **60**: 9863-9873.
- [38] Hayashi K, Takai S, Matsushima Nishiwaki R, et al. (2008). Epigallocatechin gallate reduces transforming growth factor β -stimulated HSP27 induction through the suppression of stress-activated protein kinase/c-Jun N-terminal kinase in osteoblasts. *Life Sciences*. **82**: 1012-1017.
- [39] Tabuchi M, Hayakawa S, Honda E, et al. (2013). Epigallocatechin-3-gallate suppresses transforming growth factor-beta signaling by interacting with the transforming growth factor-beta type II receptor. *World*. **3**: 100-107.
- [40] Domitrović R, Rashed K, Cvijanović O, et al. (2015). Myricitrin exhibits antioxidant, anti-inflammatory and anti-fibrotic activity in carbon tetrachloride-intoxicated mice. *Chemico-Biological Interactions*. **230**: 21-29.
- [41] Kandasamy N, Ashokkumar N. (2014). Renoprotective effect of myricetin restrains dyslipidemia and renal mesangial cell proliferation by the suppression of sterol regulatory element binding proteins in an experimental model of diabetic nephropathy. *European Journal of Pharmacology*. **743**: 53-62.
- [42] Liu HL, Jiang WB, Xie MX. (2010). Flavonoids: recent advances as anticancer drugs. *Recent Patents on Anti-Cancer Drug Discovery*. **5**: 152-164.